

**Dr. med. Henrik Behrend**

Kantonsspital St.Gallen  
ORTHO  
Switzerland

## publications

as of: 02.01.2017

### journal paper/review

Trinler U K, Baty F, Mündermann A, Fenner V, Behrend H, Jost B, Wegener R. Stair dimension affects knee kinematics and kinetics in patients with good outcome after TKA similarly as in healthy subjects. J Orthop Res 2016;.

Hochreiter B, Strahm C, Behrend H. Short-interval two-stage approach to primary total knee arthroplasty for acutely septic osteoarthritic knees. Knee Surg Sports Traumatol Arthrosc 2016;.

Ismailidis P, Kuster M S, Jost B, Giesinger K, Behrend H. Clinical outcome of increased flexion gap after total knee arthroplasty. Can controlled gap imbalance improve knee flexion?. Knee Surg Sports Traumatol Arthrosc 2016;.

Fenner V U, Behrend H, Kuster M S. Joint Mechanics After Total Knee Arthroplasty While Descending Stairs. J Arthroplasty 2016;.

Langen S, Gaber S, Zdravkovic V, Giesinger K, Jost B, Behrend H. Lateral subvastus approach with tibial tubercle osteotomy for primary total knee arthroplasty: clinical outcome and complications compared to medial parapatellar approach. Eur J Orthop Surg Traumatol 2015;.

Giesinger J M, Hamilton D F, Jost B, Behrend H, Giesinger K. WOMAC, EQ-5D and Knee Society Score Thresholds for Treatment Success After Total Knee Arthroplasty. J Arthroplasty 2015;.

Matsumoto M, Baba T, Homma Y, Kobayashi H, Ochi H, Yuasa T, Behrend H, Kaneko K. Validation study of the Forgotten Joint Score-12 as a universal patient-reported outcome measure. Eur J Orthop Surg Traumatol 2015;.

Attinger M C, Behrend H, Jost B. Complete rupture of the popliteal artery complicating high tibial osteotomy. J Orthop 2014; 11:192-6.

V F, Behrend H, MS K. Whole body gait function during stair ascent and level walking in patients following total knee arthroplasty. Int J Phys Med Rehabil 2014; 2:231-237.

Giesinger K, Behrend H. Der "Forgotten Joint Score" - Ein operiertes Gelenk im Alltag vergessen: Massstab und Ziel einer erfolgreichen Behandlung: Leading Opinions Orthopädie und Rheumatologie 1/2014; 28-30. 2014;.

Hertel P, Behrend H. BTB double bundle ACL reconstruction using a pressfit technique without fixation material. *Techniques in Orthopaedics* 2013; 2:133-140.

Giesinger J, Behrend H, Giesinger K. Association of psychological status and patient-reported physical outcome measures in joint arthroplasty: a lack of divergent validity. *Health Qual Life Outcomes* 2013; 11:1477-7525-11-64.

von der Heide N, Ebnetter L, Behrend H, Stutz G. Improvement of primary stability in ACL reconstruction by mesh augmentation of an established method of free tendon graft fixation. A biomechanical study on a porcine model. *Knee* 2013; 20:79-84.

Behrend H, Giesinger K, Giesinger J, Kuster M S. The "forgotten joint" as the ultimate goal in joint arthroplasty: validation of a new patient-reported outcome measure. *J Arthroplasty* 2012; 27:430-436.e1.

Hertel P, Behrend H. Implant-free anterior cruciate ligament reconstruction with the patella ligament and press-fit double bundle technique. *Unfallchirurg* 2010; 113(7):540-548.

MA K, Behrend H, A R, G S, MS K. Konservative oder operative Behandlung nach Ruptur des vorderen Kreuzbandes - Ein Therapiekonzept unter Berücksichtigung von Arthroseentwicklung und Funktion im Langzeitverlauf: *Rheumatologie und Traumatologie*. 2009;.

Kessler M A, Behrend H, Henz S, Rukavina A. Function, osteoarthritis and activity after ACL-rupture: 11 years follow-up results of conservative versus reconstructive treatment. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA* 2008; 16:442-8.

Behrend H, C W. Reisebericht AGA-SFA Travelling Fellowship 2006: *Arthroskopie* 2007 Mai. 20/2: 165 - 167. 2007;.

Behrend H, Stutz G, Kessler M A, Rukavina A, Giesinger K, Kuster M S. Tunnel placement in anterior cruciate ligament (ACL) reconstruction: quality control in a teaching hospital. *Knee Surg Sports Traumatol Arthrosc* 2006; 14:1159-65.

Behrend H, Bächli H. Knee ligament injuries. *Praxis* 2006; 95:1663-71.

MA K, G S, Behrend H, A R, MS K. Is ACL-Reconstruction necessary to prevent Osteoarthritis? 12 Years follow-up Results after non-operative treatment of ACL-Rupture: *British Journal of Sportsmedicine*. 2005;.

Hertel P, Behrend H, Cierpinski T, Musahl V, Widjaja G. ACL reconstruction using bone-patellar tendon-bone press-fit fixation: 10-year clinical results. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA* 2005; 13:248-55.

Behrend H, Hertel P. Results of the surgical treatment of arthro-fibrosis of the knee. *Der Unfallchirurg* 2003; 106:483-91.

## book chapter

Behrend H (2010). Erkrankungen und Verletzungen von Kniegelenk und Unterschenkel. In *Verletzungen des Kapsel-Bandapparates*: Springer-Verlag.

Behrend H (2009). Luxationen. In *Klinikmanual Chirurgie* (pp. 316-328). Heidelberg: Springer-Verlag.

M K, & Behrend H (2009). Frakturen am Körperstamm. In *Klinikmanual Chirurgie*: Heidelberg: Springer, 2009.- ISBN 978-3-540-72507-7, S. 312-315.

Behrend H (2009). Frakturen der unteren Extremität. In Klinikmanual Chirurgie (pp. 274-311). Heidelberg: Springer.

Kessler M, & Behrend H (2009). Frakturen der oberen Extremität. In Klinikmanual Chirurgie (pp. 250-273). Heidelberg: Springer.

Behrend H (2009). Weichteilverletzungen. In Klinikmanual Chirurgie (pp. 329-341). Heidelberg: Springer.