



Dr. Walter Brunner

Allgemein-, Viszeral-, Endokrin- und Transplantationschirurgie · Dept. II

Kontakt

Dr. Walter Brunner
Switzerland

Walter.Brunner@kssg.ch

Bereiche

Allgemein-, Viszeral-, Endokrin- und Transplantationschirurgie

Publikationen (25)

Tschann P, Brock T, Weigl M, Clemens P, Brunner W, Attenberger C, Königsrainer I, Emmanuel K, Jäger T. Tumour, narrow pelvis and surgery specific factors for total mesorectal excision quality and morbidity following rectal cancer resection. *BJS Open* 2023; 7

Bischofberger S, Adshead F, Moore K, Kocaman M, Casali G, Tong C, Roy S, Collins M, Brunner W. Assessing the environmental impact of an anastomotic leak care pathway. *Surg Open Sci* 2023; 14:81-86.

Pietrogiiovanna L, Janczak J, Pfeifer N, Strahm R, Brunner W. Incarcerated femoral hernia in women – A critical view on approach options. *Int J Surg Case Rep* 2023; 106:108149.

Tschann P, Weigl M, Brockamp T, Frick J, Sturm O, Presl J, Jäger T, Weitzendorfer M, Schredl P, Clemens P, Eiter H, Szeverinski P, Attenberger C, Tschann V, Brunner W, de Vries A, Emmanuel K, Königsrainer I. Identification of Risk Factors for Sexual Dysfunction after Multimodal Therapy of Locally Advanced Rectal Cancer and Their Impact on Quality of Life: A Single-Center Trial. *Cancers (Basel)* 2022; 14:5796.

Widmann B, Almarie B, Warschkow R, Beutner U, Weitzendorfer M, Schmid M, Ukegini K, Brunner W. Advanced Lymph Node Staging with Ex Vivo Intraarterial Indigo Carmine Injection after Transanal Total Mesorectal Excision for Rectal Cancer: A Retrospective Cohort Study. *Dis Colon Rectum* 2021; 65:1015-1024.

Ammann Y, Widmann B, Sparn M, Warschkow R, Weitzendorfer M, Brunner W. Prophylactic Funnel Mesh to Prevent Parastomal Hernia in Permanent End Colostomy: A Retrospective Cohort Study. *Colorectal Dis* 2021; 23:2627-2636.

Köckerling F, Zarras K, Reinpold W, Lorenz R, Niebuhr H, Adolf D, Mayer F, Fortelny R, Brunner W, Weyhe D. Treatment of small (< 2 cm) umbilical hernias: guidelines and current trends from the Herniated Registry. *Hernia* 2020; 25:605-617.

Köckerling F, Zarras K, Reinpold W, Lorenz R, Niebuhr H, Adolf D, Fortelny R, Mayer F, Brunner W, Weyhe D. Assessment of potential influencing factors on the outcome in small (< 2 cm) umbilical hernia repair: a registry-based multivariable analysis of 31,965 patients. *Hernia* 2020

TaTMe Guidance Group representing the ESCP in collaboration with the ASCRS, ACPGBI, ECCO, EAES, ESSO, CSCRS, CNSCRS, CSLES, CSSANZ, JSES, SACP, SAGES, SBCP, Swiss-MIS, Brunner W. International expert consensus guidance on indications, implementation and quality measures for transanal total mesorectal excision. *Colorectal Dis* 2020; 22:749-755.

Brunner W. Onkologische und chirurgische Ergebnisse der kolorektalen roboterassistierten Chirurgie – Wie gehen wir mit Lokalrezidivraten um?. *coloproctology* 2020; 42:47-48.

Konschake M, Zwierzina M, Moriggl B, Függer R, Mayer F, Brunner W, Schmid T, Chen D, Fortelny R. The inguinal region revisited: the surgical point of view : An anatomical-surgical mapping and sonographic approach regarding postoperative chronic groin pain following open hernia repair. *Hernia* 2019; 24:883-894.

Roodbeen S, International TaTME Registry Collaborative, Hompes R, Bemelman W, Tekkis P, Moran B, Ris F, Penna M, van Dieren S, de Lacy F, Brunner W. Predictive Factors and Risk Model for Positive Circumferential Resection Margin Rate After Transanal Total Mesorectal Excision in 2653 Patients With Rectal Cancer. *Ann Surg* 2019; 270:884-891.

Penna M, International TaTME Registry Collaborative, Tekkis P, Mortensen N, Hanna G, Moran B, Warusavitarne J, Austin R, Wynn G, Arnold S, Hompes R, Brunner W. Incidence and Risk Factors for Anastomotic Failure in 1594 Patients Treated by Transanal Total Mesorectal Excision: Results From the International TaTME Registry. <i>Ann Surg</i> 2019; 269:700–711.
Wilhelm A, Galata C, Beutner U, Schmied B, Warschkow R, Steffen T, Brunner W, Post S, Marti L. Duodenal localization is a negative predictor of survival after small bowel adenocarcinoma resection: A population-based, propensity score-matched analysis. <i>J Surg Oncol</i> 2017
Marti L, Sulz M, Brunner W, Borovicka J, Wolff K, Pipitone N, Hetzer F, Beutner U, Galata C, Maurus C. Percutaneous tibial nerve stimulation (pTNS): success rate and the role of rectal capacity. <i>Int J Colorectal Dis</i> 2017; 32:789–796.
Warschkow R, Ebinger S, Brunner W, Schmied B, Marti L. Survival after Abdominoperineal and Sphincter-Preserving Resection in Nonmetastatic Rectal Cancer: A Population-Based Time-Trend and Propensity Score-Matched SEER Analysis. <i>Gastroenterol Res Pract</i> 2017; 2017:6058907.
Weiss H, Hell T, Muratore A, Weiss M, Stanger O, Klaus A, Mittermair C, Dixon A, Gash K, Bulut O, Morales Conde S, Sietes C, Brunner W, Boni L, Vestweber B, Vestweber K, Zorron R. ECSPECT prospective multicentre registry for single-port laparoscopic colorectal procedures. <i>Br J Surg</i> 2016
Steinemann D, Vorburger S, Spalinger R, Scheiwiller A, Nocito A, Keerl A, Brunner W, Adamina M, Zerz A, Lamm S. Single-Incision and Natural Orifice Transluminal Endoscopic Surgery in Switzerland. <i>World J Surg</i> 2016
Hummel B, Schmied B, Widmann B, Brunner W, Zadnikar M, Warschkow R, Hetzer F, Bischofberger S, Hardt J, Marti L. New kid on the block: perineal stapled prolapse resection (PSP) is it worthwhile in the long-term?. <i>Langenbecks Arch Surg</i> 2016; 401:519–29.
Kruse C, Tarantino I, Schmied B, Sulz M, Marti L, Brunner W, Luethi C, Warschkow R, Mueller S, Beutner U. Does caffeine reduce postoperative bowel paralysis after elective laparoscopic colectomy? (CaCo trial): study protocol for a randomized controlled trial. <i>Trials</i> 2016; 17:186.
Brunner W, Widmann B, Marti L, Tarantino I, Schmied B, Warschkow R. Predictors for regional lymph node metastasis in T1 rectal cancer: a population-based SEER analysis. <i>Surg Endosc</i> 2016; 30:4405–15.
Sawatzki M, Brunner W, Neff-Schöb A, Meyenberger C, Borovicka J. Chronische Nausea und Emesis. <i>Swiss Medical Forum</i> 2015; 15:1039–1040.
Brunner W, Rossetti A, Vines L, Kalak N, Bischofberger S. Anastomotic leakage after laparoscopic single-port sigmoid resection: combined transanal and transabdominal minimal invasive management. <i>Surg Endosc</i> 2015; 29:3803–5.
Aigner F, Morales-Conde S, Zorron R, Sileri P, Boni L, de Graaf E, Scherer R, Shamiyeh A, Klaus A, Tse C, Brunner W, Panis Y, Herbst F, Hörmann R, Fritsch H, Pratschke J, D'Hoore A, Brenner E, Williams N, Biebl M, The TAMIS TME collaboration group, Weiss H, Mittermair R. Anatomical considerations for transanal minimal-invasive surgery: the caudal to cephalic approach. <i>Colorectal Dis</i> 2015; 17:O47–O53.
Weiss H, Brunner W, Biebl M, Schirnhöfer J, Pimpl K, Mittermair C, Obrist C, Brunner E, Hell T. Wound complications in 1145 consecutive transumbilical single-incision laparoscopic procedures. <i>Ann Surg</i> 2014; 259:89–95.

Projekte (3)

Low Anterior Resection Syndrome After Rectum Resection for Rectal Carcinoma. (LARS-Austria)

Klinische Forschung - 01.01.2022 - 30.08.2024

Laufend

Randomisierte klinische Studie zum Vergleich der Vollständigkeit der adjuvanten Chemotherapie nach früher vs. später Stomarückverlagerung bei Patienten mit Rektumkarzinom nach tiefer anteriorer Resektion

Klinische Forschung - 24.09.2018 - 30.11.2020

Abgeschlossen

CaCo: Does caffeine intake reduce postoperative bowel paralysis after elective laparoscopic colectomy? - A randomized placebo controlled trial

Klinische Forschung - 01.08.2015 - 31.07.2022

Automatisch geschlossen